



Treatment guidelines for P.falciparum and Mixed



ASMQ

Weight	Age	Recommended dose	Day 1	Day 2	Day 3
5 – 8 kg	6 – 11 months	One Tablet 25/50 mg daily for 3 days			
9 – 17 kg	1 – 6 years	Two Tablets 25/50 mg daily for 3 days			
18 – 29 kg	7 – 12 years	One Tablet 100/200 mg daily for 3 days			
≥ 30 kg	≥ 13 years	Two Tablets 100/200 mg daily for 3 days			


ASMQ Dosing:

- **Artesunate + Mefloquine** dosages are available in **25/50 mg** and **100/200 mg** sized tablets
- Dosage and number of pills used are **dependent on weight**, if the weight is not known, use age ranges

Do not use ASMQ for:

- 1 Patients who have received **Mefloquine** in the **last 4 weeks**
- 2 Patients with a **history of seizures or epilepsy**
- 3 **Pregnant women** in their **first trimester**

If patient vomits after treatment of ASMQ



Primaquine

Administer single low dose Primaquine for P.f. and mix only on the first day of ASMQ treatment


Weight	Public Health Facilities		VMWs	
	# Tablets		# Tablets	
20kg - 49kg	1 tablet of 7.5mg		1 tablet of 7.5mg	
≥ 50kg	1 tablet of 15mg		2 tablets of 7.5mg	

Do not treat pregnant and breastfeeding women

Tell patient to always take with food!

If patient vomits, no Primaquine shall be given to patient after the vomit.

Positive



or

Malaria (P.falciparum)

Malaria (Mix)